

Approved:
CEO: /Ch. Marinov/
Date: 22.06.23

Outgoing Ref. No. 14.03.2024
(filled in by CARBONSAFE)

Individual strategy for management of used areas in the farm

Data of the operator:

Natural person ☐	Legal entity ☒
DATA OF THE NATURAL PERSON (filled in by Natural persons only)	
Name:	
BULSTAT:	
Telephone:	
e-mail:	
DATA OF THE LEGAL ENTITY (filled in by Legal entities only)	
Name:	Kris-Mari-2008 OOD
UIC:	200359382
Telephone:	0878688060
e-mail:	krismari2008@gmail.com

Table 1:

No. in row	CROPS GROWN IN THE FARM	ID No. of the plot (by Restored Property Map/Cadastral Map/Agricultural Land Block/Farming Block)	Area ha	EXISTING SITUATION (Short description of the applied technology and types of tillage/treatment - visual agronomic assessment)			
				1 ☐	6 X	11 ☐	16 X
1	Winter soft wheat		172,001	2 ☐	7 X	12 ☐	17 ☐
				3 X	8 X	13 ☐	18 ☐
				4 ☐	9 ☐	14 X	19 ☐
				5 ☐	10 ☐	15 ☐	20 ☐
2	Winter rapeseed		87,768	1 ☐	6 X	11 ☐	16 X
				2 ☐	7 X	12 ☐	17 ☐
				3 X	8 X	13 ☐	18 ☐
				4 ☐	9 ☐	14 X	19 ☐
				5 ☐	10 ☐	15 ☐	20 ☐
3	Sunflower		138,66	1 ☐	6 X	11 ☐	16 X
				2 ☐	7 X	12 ☐	17 ☐
				3 X	8 X	13 ☐	18 ☐
				4 ☐	9 ☐	14 X	19 ☐
				5 ☐	10 ☐	15 ☐	20 ☐
4	Grain maize		88,439	1 ☐	6 X	11 ☐	16 X
				2 ☐	7 X	12 ☐	17 ☐
				3 X	8 X	13 ☐	18 ☐
				4 ☐	9 ☐	14 X	19 ☐
				5 ☐	10 ☐	15 ☐	20 ☐

LEGEND	
☐ 1. Conservation tillage / no till;	☐ 13. Stripes tillage;
☐ 2. Treatment of beds;	☐ 14. Pasture and/or crop rotation and crop rotation management;
☐ 3. Minimal treatments;	

☐ 4. Biological agriculture; ☐ 5. Integrated production; ☐ 6. Precision agriculture; ☐ 7. Diversification of crops; ☐ 8. Fertilization with microbial fertilizers; ☐ 9. Green fertilization (Sideration); ☐ 10. Cultivation of nitrogen-fixing crops; ☐ 11. Mulching treatment; ☐ 12. Use of organic/natural pesticides;	☐ 15. Grass weeding of the rows in perennial crops and vineyards; ☐ 16. Joint cultivation of more than one agricultural crop; ☐ 17. Implementation of agricultural belts; ☐ 18. Improvement measures in permanently grassed areas; ☐ 19. OTHER NOT LISTED HERE; ☐ 20. NONE OF THE ABOVE APPLIES**.
---	---

ADDITIONAL INFORMATION: (Please describe cultivation methods and practices, including tillage, currently being carried out on the cultivated areas and not listed above:)

.....

Table 2:

No. in row	CROPS GROWN IN THE FARM	ID No. of the plot (by Restored Property Map/Cadastral Map/Agricultural Land Block/Farming Block)	Area ha	INDIVIDUAL STRATEGY FOR MANAGEMENT OF THE USED AREAS IN THE FARM			
1	Sunflower		172,001	☐ 1 ☐ 2 ☒ 3 ☐ 4 ☒ 5	☒ 6 ☒ 7 ☒ 8 ☐ 9 ☐ 10	☐ 11 ☐ 12 ☐ 13 ☒ 14 ☐ 15	☒ 16 ☐ 17 ☐ 18 ☐ 19 ☐ 20
2	Durum wheat		314,6111	☒ 1 ☐ 2 ☒ 3 ☐ 4 ☒ 5	☒ 6 ☒ 7 ☒ 8 ☐ 9 ☐ 10	☐ 11 ☐ 12 ☐ 13 ☒ 14 ☐ 15	☒ 16 ☐ 17 ☐ 18 ☐ 19 ☐ 20

LEGEND	
☐ 1. Conservation tillage / no till; ☐ 2. Treatment of beds; ☐ 3. Minimal treatments; ☐ 4. Biological agriculture; ☐ 5. Integrated production; ☐ 6. Precision agriculture; ☐ 7. Diversification of crops; ☐ 8. Fertilization with microbial fertilizers; ☐ 9. Green fertilization (Sideration); ☐ 10. Cultivation of nitrogen-fixing crops; ☐ 11. Mulching treatment; ☐ 12. Use of organic/natural pesticides;	☐ 13. Stripes tillage; ☐ 14. Pasture and/or crop rotation and crop rotation management; ☐ 15. Grass weeding of the rows in perennial crops and vineyards; ☐ 16. Joint cultivation of more than one agricultural crop; ☐ 17. Implementation of agricultural belts; ☐ 18. Improvement measures in permanently grassed areas; ☐ 19. Other not listed here; ☐ 20. None of the above applies**.

ADDITIONAL INFORMATION: (Please describe cultivation methods and practices, including tillage, to be carried out on cultivated areas not listed above:)

.....

No. from the legend	Designation	Notes
---------------------	-------------	-------

3	Minimal treatments: Opti-till or Minimum-till - They are characterized by minimal, partial, merged soil treatments, one, two or three in number depending on the type of soil and the specific climatic conditions. This is a technology in which deep soil treatments are replaced by the construction of a biological soil eco-system. In it, microorganisms, roots and other soil fauna take over the functions of processing and balancing nutrients. Minimal treatments preserve carbon in the soil, protect it from water and wind erosion, prevent compaction, improve soil fertility. Soil fertility (nutrients and water) is regulated by managing the soil cover from the previous crop, partial soil treatments and weed control.	 Sunflower Durum wheat
5	Integrated production: Integrated production is a certified quality system for the production of crops that supports environmental protection through integrated pest management (IPM) and reducing the use of PPPs (plant protection products). Unlike organic farming, which excludes the use of PPPs or mineral fertilizers, in integrated production they can be applied, but under certain conditions. Integrated production makes use of advances in technology in the cultivation and protection of crops and combines different methods and means of pest management. Each farmer can apply integrated pest management by including a number of preventive measures to limit the spread of pests – such as crop rotation, use of appropriate agricultural machinery, balanced fertilization and watering, sanitary and hygienic measures, protection of beneficial organisms, resistant / tolerant plant varieties and standard/certified seeds and planting material, etc. Systematic monitoring of pests is carried out. When controlling them, preference is given to biological, physical, biotechnical and other non-chemical means, as well as to low-risk PPPs.	 Sunflower Durum wheat
6	Precision agriculture: Precision agriculture or farming is based on the use of a wide range of technologies that allow the collection of data from treatments, monitoring and analysis of crop development, as areas are adequately treated to increase efficiency. This management system is based on decision making, based on variable characteristics and obtaining maximum yields, according to the specifics of the site. The main benefits are associated with reduced use of water, fertilizers and pesticides, depending on specific data on conditions and a set of necessary agrotechnical measures. Precision agriculture should also be developed to support the development of precision technologies for sustainable agriculture in a clean and secure environment.	 Sunflower Durum wheat
7	Diversification of crops: • Crop diversification - is the cultivation of several different crops on the farm in order to avoid monocultural production, which can lead to a decrease in soil fertility, increasing problems with protection from diseases, pests and weeds, which in turn is a prerequisite for greater levels of fertilizer input and plant protection products. On a farm with arable land between 10 ha and 30 ha (inclusive) the farmer must provide at least 2 different crops. The main crop should not cover more than 75% of the arable land.	 Sunflower Durum wheat

	A holding with arable land above 30 ha should have at least 3 different crops. The main crop should not cover more than 75% of the arable land, and the two main crops should not be more than 95% of arable land. The diversification requirement shall not apply where: - More than 75 % of arable land is used for the production of grasses or other herbaceous forage, sown with legumes, set aside or combined with these uses and, if it is permanent grassland, for the production of grasses or other herbaceous forage or for the production of crops underwater. - Arable land on the farm is up to 10 ha.	
8	Fertilization with microbial fertilizers: Microbials or so-called 'live fertilizers' means substances containing live micro-organisms which, when applied to seeds, plant surfaces or soil, colonize the rhizosphere or plant interior and promote growth by increasing the supply or availability of essential nutrients to the host plant. These fertilizers add nutrients through the natural processes of nitrogen fixation, dissolution of phosphorus and stimulation of plant growth by the synthesis of growth-promoting substances. It is expected that in the future, microbial fertilizers will significantly reduce the use of chemical fertilizers and pesticides. Benefits: Reduced use of mineral fertilizers. Improved accessibility of nutrients. Improvement of biological activity and soil fertility.	 Sunflower Durum wheat
14	Pasture and/or crop rotation and crop rotation management: Integration of different agricultural crops in agricultural areas, including cover crops. Crop rotation is a scientifically based alternation of crops in time and place on a certain area of a farm. Time rotation consists in changing crops in successive years on the same field. The rotation by place consists in the successive passage of each crop through all the fields. The rotation must meet the requirements of modern agronomic science, be rational, provide an economically advantageous crop structure for the farm, be in accordance with the ecological requirements of the crops and meet the terrain and relief conditions. Long-term cultivation of the same crop in one place causes a gradual decrease in soil fertility, an increase in the concentration of diseases and enemies. This can be avoided if crops are grown in crop rotation. In this way, biological factors are used most effectively to maintain and increase soil fertility. In order to ensure the rotation of crops in time and place, it is necessary to divide the total sowing rotation area into separate fields (most often 4-6). Plots of cultivated area that are occupied by one or several crops (when collective fields are formed) and that have approximately the same dimensions are called crop rotation fields. Benefits: Reduced use of pesticides and fertilizers through the inclusion of leguminous crops. Reduction of threats from wind and water erosion of the areas occupied by roof vegetation.	 Sunflower Durum wheat

16	Joint cultivation of more than one agricultural crop: Inter-row cropping or parallel growing of more than one crop on the same area. Applied mainly by smaller farms, but the selected agricultural crops must be adapted to the climatic characteristics of the area. Benefits: Increases the productivity of the land because it saves space. Reducing the impact of weeds and pests. Improving soil nutrient content in growing legumes. Disadvantages: Limitations in technical harvesting equipment. The choice of crops is based on the agro-ecological characteristics of the area. • Nutrient management and fertilization practices. Nutrient management (NPK) - nutrient balance. The introduction of mineral fertilizers must be carried out taking into account the needs of the planned crop, which will be grown based on analyzes carried out for the presence of a number of trace elements and residues of nitrogen, phosphorus and potassium in the soil. When calculating the fertilization rate, a number of indicators are taken into account: soil type and mineral composition, culture-predecessor, ratio between N/P/K. The quantities of mineral fertilizers and nutrients are imported only on the basis of a precisely calculated fertilizer rate. The practice of applying only nitrogen fertilizers is categorically denied. The required minimum balance between N/P/K calculated for the specific crop and the specific field on which it will be grown is sought.	 Sunflower Durum wheat
-----------	---	---

Date 14.03.2024

Agronomist: H. Nikolov