

Approved:
CEO: /Ch. Marinov/
Date: 22.06.23

Outgoing Ref. No. 29.04.2024
(filled in by CARBONSAFE)

Individual strategy for management of used areas in the farm

Data of the operator:

Natural person ☐	Legal entity ☒
DATA OF THE NATURAL PERSON (filled in by Natural persons only)	
Name:	
BULSTAT:	
Telephone:	
e-mail:	
DATA OF THE LEGAL ENTITY (filled in by Legal entities only)	
Name:	PROIZVODITEL YOTIN EOOD
UIC:	207021636
Telephone:	+359886555249
e-mail:	proizvoditel.yotin@abv.bg

Table 1:

No. in row	CROPS GROWN IN THE FARM	ID No. of the plot (by Restored Property Map/Cadastral Map/Agricultural Land Block/Farming Block)	Area ha	EXISTING SITUATION (Short description of the applied technology and types of tillage/treatment - visual agronomic assessment)			
				1	2	3	4
1	Sunflower		63,877	1 ☒ 2 ☐ 3 ☒ 4 ☒ 5 ☒	6 ☒ 7 ☒ 8 ☒ 9 ☐ 10 ☐	11 ☐ 12 ☒ 13 ☒ 14 ☐ 15 ☐	16 ☐ 17 ☒ 18 ☐ 19 ☐ 20 ☐
2	Wheat		20,466	1 ☒ 2 ☐ 3 ☒ 4 ☒ 5 ☒	6 ☒ 7 ☒ 8 ☒ 9 ☐ 10 ☐	11 ☐ 12 ☒ 13 ☐ 14 ☐ 15 ☐	16 ☐ 17 ☒ 18 ☐ 19 ☐ 20 ☐
3	Alfalfa		1,752	1 ☒ 2 ☐ 3 ☒ 4 ☒ 5 ☒	6 ☒ 7 ☒ 8 ☒ 9 ☒ 10 ☒	11 ☐ 12 ☒ 13 ☐ 14 ☒ 15 ☐	16 ☒ 17 ☒ 18 ☐ 19 ☐ 20 ☐
4	Spelt		4,740	1 ☒ 2 ☐ 3 ☒ 4 ☒ 5 ☒	6 ☒ 7 ☒ 8 ☒ 9 ☐ 10 ☐	11 ☐ 12 ☒ 13 ☐ 14 ☐ 15 ☐	16 ☐ 17 ☒ 18 ☐ 19 ☐ 20 ☐

LEGEND	
☐ 1. Conservation tillage / no till; ☐ 2. Treatment of beds; ☐ 3. Minimal treatments;	☐ 13. Stripes tillage; ☐ 14. Pasture and/or crop rotation and crop rotation management;

☐ 4. Biological agriculture; ☐ 5. Integrated production; ☐ 6. Precision agriculture; ☐ 7. Diversification of crops; ☐ 8. Fertilization with microbial fertilizers; ☐ 9. Green fertilization (Sideration); ☐ 10. Cultivation of nitrogen-fixing crops; ☐ 11. Mulching treatment; ☐ 12. Use of organic/natural pesticides;	☐ 15. Grass weeding of the rows in perennial crops and vineyards; ☐ 16. Joint cultivation of more than one agricultural crop; ☐ 17. Implementation of agricultural belts; ☐ 18. Improvement measures in permanently grassed areas; ☐ 19. OTHER NOT LISTED HERE; ☐ 20. NONE OF THE ABOVE APPLIES**.
---	---

ADDITIONAL INFORMATION: (Please describe cultivation methods and practices, including tillage, currently being carried out on the cultivated areas and not listed above:)

.....

Table 2:

No. in row	CROPS GROWN IN THE FARM	ID No. of the plot (by Restored Property Map/Cadastral Map/Agricultural Land Block/Farming Block	Area ha	INDIVIDUAL STRATEGY FOR MANAGEMENT OF THE USED AREAS IN THE FARM			
1	Sunflower		6,4625	1 ☒ 2 ☐ 3 ☒ 4 ☒ 5 ☒	6 ☒ 7 ☒ 8 ☒ 9 ☐ 10 ☐	11 ☐ 12 ☒ 13 ☒ 14 ☐ 15 ☐	16 ☐ 17 ☒ 18 ☐ 19 ☐ 20 ☐
2	Wheat		59,3233	1 ☒ 2 ☐ 3 ☒ 4 ☒ 5 ☒	6 ☒ 7 ☒ 8 ☒ 9 ☐ 10 ☐	11 ☐ 12 ☒ 13 ☐ 14 ☐ 15 ☐	16 ☐ 17 ☒ 18 ☐ 19 ☐ 20 ☐
3	Triticale		7,2185	1 ☒ 2 ☐ 3 ☒ 4 ☒ 5 ☒	6 ☒ 7 ☒ 8 ☒ 9 ☐ 10 ☐	11 ☐ 12 ☒ 13 ☐ 14 ☐ 15 ☐	16 ☐ 17 ☒ 18 ☐ 19 ☐ 20 ☐
4	Barley		17,8022	1 ☒ 2 ☐ 3 ☒ 4 ☒ 5 ☒	6 ☒ 7 ☒ 8 ☒ 9 ☐ 10 ☐	11 ☐ 12 ☒ 13 ☐ 14 ☐ 15 ☐	16 ☐ 17 ☒ 18 ☐ 19 ☐ 20 ☐

LEGEND

- | | |
|---|---|
| ☐ 1. Conservation tillage / no till;
☐ 2. Treatment of beds;
☐ 3. Minimal treatments;
☐ 4. Biological agriculture;
☐ 5. Integrated production;
☐ 6. Precision agriculture;
☐ 7. Diversification of crops;
☐ 8. Fertilization with microbial fertilizers;
☐ 9. Green fertilization (Sideration);
☐ 10. Cultivation of nitrogen-fixing crops;
☐ 11. Mulching treatment;
☐ 12. Use of organic/natural pesticides; | ☐ 13. Stripes tillage;
☐ 14. Pasture and/or crop rotation and crop rotation management;
☐ 15. Grass weeding of the rows in perennial crops and vineyards;
☐ 16. Joint cultivation of more than one agricultural crop;
☐ 17. Implementation of agricultural belts;
☐ 18. Improvement measures in permanently grassed areas;
☐ 19. Other not listed here;
☐ 20. None of the above applies**. |
|---|---|

ADDITIONAL INFORMATION: (Please describe cultivation methods and practices, including tillage, to be carried out on cultivated areas not listed above:)

.....
.....

No. from the legend	Designation	Notes
1	DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues at periods of 2-3 years depending on weather conditions. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from June 1 to October 31, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map. Conservation tillage /No-till: Conservation tillage is any cultivation or crop growing system that, after sowing, leaves at least 30% (or more) of crop residue on the soil surface after sowing to reduce water erosion. Any soil cultivation system that maintains at least 90-100 kg/ha of crop residue (stubble) during a critical wind-erosion period is considered to be conservation treatment. Effective conservation agriculture is based on three basic principles: • minimal mechanical disturbances of the soil (i.e., without tillage) by direct placement of seeds and / or fertilizer. • permanent soil organic cover, with at least 30% with crop residues and/or cover crops. • diversification of crops grown successively and/or in association. By reducing treatments, they can save between 30 and 40% of the labor time and petroleum fuels needed for mechanized activities compared to conventional farming. This would have a positive impact on the emissions of gases - pollutants of the atmosphere. Conservation agriculture also has the following advantages: • A sustainable agricultural system, (not only conservation), which improves the quality of	 Sunflower Wheat Triticale Barley

	natural resources, increases soil biodiversity, (flora and fauna, including wildlife) and at the same time does not confront the desire to obtain higher yields. • Uncultivated (unplugged) fields act as a landfill for carbon dioxide (CO₂) and applied globally, can be important in controlling atmospheric pollution in general, and in particular in influencing global warming. • Soils treated through conservation practices have better infiltration ratio and reduced surface runoff, which significantly reduces erosion. The practices for minimum mechanical disturbances of soils can be divided into two more general groups – with minimal treatments and without tillage (direct sowing). No tillage - Direct sowing in living roof vegetation or mulch. Direct sowing without plowing in the previously available roof vegetation – living or destroyed, e.g., mulch. Benefits: Reduced costs associated with pre-sowing treatments not carried out. Reduction of threats from wind- and water-caused soil erosion. Reduction of soil compaction and increased bioactivity. Reducing evaporation and excessive soil moisture loss, often critical for crop development. Less development of weeds in the long term, leading to less use of herbicides. Disadvantages: Difficulties for effective control of weeds, cover vegetation and mulch. Increased costs associated with mechanical weed control. Need for special machines, such as a seeder for direct sowing and other specialized machines.	
3	DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map.	 Sunflower Wheat Triticale Barley


	Minimal treatments: Opti-till or Minimum-till - They are characterized by minimal, partial, merged soil treatments, one, two or three in number depending on the type of soil and the specific climatic conditions. This is a technology in which deep soil treatments are replaced by the construction of a biological soil eco-system. In it, microorganisms, roots and other soil fauna take over the functions of processing and balancing nutrients. Minimal treatments preserve carbon in the soil, protect it from water and wind erosion, prevent compaction, improve soil fertility. Soil fertility (nutrients and water) is regulated by managing the soil cover from the previous crop, partial soil treatments and weed control.	
4	DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map. Biological agriculture: Organic farming is an aggregate agricultural management and food production system that combines best environmental practices, maintains a high level of biodiversity, conserves natural resources, applies highly humane animal treatment and welfare standards and production methods tailored to the preferences of some consumers for products, produced using natural substances and processes. For compliance with the requirements of the applicable European legislation, annual control is carried out and the production is subject to certification after the transition periods have passed. Organic farming is a production system that prevents or completely excludes the use of synthetic fertilizers, pesticides, growth regulators and additives to animal feed, and in which crop rotations, plant residues, manure, green fertilization and biological plant protection are relied on to maintain and improve the soil diet.	 Sunflower Wheat Triticale Barley

5	DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map. Integrated production: Integrated production is a certified quality system for the production of crops that supports environmental protection through integrated pest management (IPM) and reducing the use of PPPs (plant protection products). Unlike organic farming, which excludes the use of PPPs or mineral fertilizers, in integrated production they can be applied, but under certain conditions. Integrated production makes use of advances in technology in the cultivation and protection of crops and combines different methods and means of pest management. Each farmer can apply integrated pest management by including a number of preventive measures to limit the spread of pests – such as crop rotation, use of appropriate agricultural machinery, balanced fertilization and watering, sanitary and hygienic measures, protection of beneficial organisms, resistant / tolerant plant varieties and standard/certified seeds and planting material, etc. Systematic monitoring of pests is carried out. When controlling them, preference is given to biological, physical, biotechnical and other non-chemical means, as well as to low-risk PPPs.	 Sunflower Wheat Triticale Barley
6	DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to	 Sunflower Wheat

	the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map. Precision agriculture: Precision agriculture or farming is based on the use of a wide range of technologies that allow the collection of data from treatments, monitoring and analysis of crop development, as areas are adequately treated to increase efficiency. This management system is based on decision making, based on variable characteristics and obtaining maximum yields, according to the specifics of the site. The main benefits are associated with reduced use of water, fertilizers and pesticides, depending on specific data on conditions and a set of necessary agrotechnical measures. Precision agriculture should also be developed to support the development of precision technologies for sustainable agriculture in a clean and secure environment.	Triticale Barley
7	DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map. DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues. • Inadmissible soil treatments: plowing and overturning of soil layers.	 Sunflower Wheat Triticale Barley

	GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map. Diversification of crops: • Crop diversification - is the cultivation of several different crops on the farm in order to avoid monocultural production, which can lead to a decrease in soil fertility, increasing problems with protection from diseases, pests and weeds, which in turn is a prerequisite for greater levels of fertilizer input and plant protection products. On a farm with arable land between 10 ha and 30 ha (inclusive) the farmer must provide at least 2 different crops. The main crop should not cover more than 75% of the arable land. A holding with arable land above 30 ha should have at least 3 different crops. The main crop should not cover more than 75% of the arable land, and the two main crops should not be more than 95% of arable land. The diversification requirement shall not apply where: - More than 75 % of arable land is used for the production of grasses or other herbaceous forage, sown with legumes, set aside or combined with these uses and, if it is permanent grassland, for the production of grasses or other herbaceous forage or for the production of crops underwater. - Arable land on the farm is up to 10 ha.	
8	DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to	 Sunflower Wheat

	the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map. Fertilization with microbial fertilizers: Microbials or so-called 'live fertilizers' means substances containing live micro-organisms which, when applied to seeds, plant surfaces or soil, colonies the rhizosphere or plant interior and promote growth by increasing the supply or availability of essential nutrients to the host plant. These fertilizers add nutrients through the natural processes of nitrogen fixation, dissolution of phosphorus and stimulation of plant growth by the synthesis of growth-promoting substances. It is expected that in the future, microbial fertilizers will significantly reduce the use of chemical fertilizers and pesticides. Benefits: Reduced use of mineral fertilizers. Improved accessibility of nutrients. Improvement of biological activity and soil fertility.	Triticale Barley
12	DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map. Use of organic/natural pesticides: Use of plant protection products (PPPs) whose active substances are based on plant extracts (e.g., pyrethrin-based PPPs extracted from chrysanthemum flowers), extracts based on extracted essential oils or extracts from recycled and processed food industry waste (citrus, fruit, vegetables, woody, shrub and flower	 Sunflower Wheat Triticale Barley

	species). Benefits: Reduction or absence of contamination of soils, water or production by chemical synthetic PPPs. Reduction of PPP risks to human health. Allowed for use as a method of plant protection in organic production.	
13	DIRECTIONS: • Recommended soil treatments: strip vertical treatments in the sowing lines. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the technological map. Stripes tillage: Strip-Till (strip tillage) – in this treatment the soil remains untilled from the harvesting of the previous crop until sowing the next one except for strips 1/3 of the row width.	 Sunflower
17	DIRECTIONS: • Recommended soil treatments: vertical treatments to mix plant residues. • Inadmissible soil treatments: plowing and overturning of soil layers. GUIDELINES: • Summer period from 01 June to 31 October, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. • Winter period from November 1 to February 15, the areas must be occupied with a minimum soil cover, plowing and overturning of soil layers is not allowed, except for a period within 14 days in which the soil can be processed according to the needs for the next crop, including sowing. DUTIES: • Every single event, including soil treatments, is marked with the relevant deadlines in the	 Sunflower Wheat Triticale Barley

	technological map. Implementation of agricultural belts: Strip agriculture is the division of the slope surface into strips of a certain width located along the horizontals of the terrain, or across the slope of the slope, on which various crops are grown such as: cereals, field crops, perennial grasses, etc. The essence of this anti-erosion method consists in successive alternation along the length of the slope of belts of trench crops with belts of crops with a fused surface. This achieves both a reduction in erosion processes and an increase in soil moisture on the slopes. The intercropped belts are a barrier that reduces the rate of surface water runoff and soil erosion, but also serve as a filter to retain entrained sediment from the upper trenched crop belt.	
--	--	--

Date 29.04.2024

Agronomist:
(L. Aleksandrova, LA0424CS)